



## King's Research Portal

DOI:

[10.1183/13993003.00837-2019](https://doi.org/10.1183/13993003.00837-2019)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Walsh, J. A., Maddocks, M., & Man, W. D. C. (2019). Supplemental oxygen during exercise training in COPD: full of hot air? *The European respiratory journal*, 53(5). <https://doi.org/10.1183/13993003.00837-2019>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## **Supplementary oxygen during exercise training in COPD - full of hot air?**

Mrs Jessica A Walsh <sup>1</sup>, Dr Matthew Maddocks <sup>2</sup>, Dr William D-C Man <sup>1,3</sup>

### **Affiliations:**

1. Harefield Respiratory Research Group, Royal Brompton & Harefield NHS Foundation Trust, UK
2. Cicely Saunders Institute of Palliative Care, Policy & Rehabilitation, King's College London, UK
3. National Heart and Lung Institute, Imperial College London, UK

### **Correspondence:**

Dr William D-C Man. Harefield Respiratory Research Group, Department of Respiratory Medicine, Harefield Hospital, Hill End Rd, Harefield, Uxbridge UB9 6JH, UK.

Email: [w.man@rbht.nhs.uk](mailto:w.man@rbht.nhs.uk)

Long term oxygen therapy is widely accepted as a treatment for severe hypoxaemia in patients with chronic respiratory disorders, based on survival findings from the UK Medical Research Council study[1] and the US Nocturnal Oxygen Therapy Trial[2]. The data from these landmark studies have driven the underlying rationale that correcting hypoxaemia is beneficial, and have led to the prescription of oxygen for other indications, including ambulatory oxygen for patients with exercise-induced oxygen desaturation (EID), nocturnal oxygen therapy for patients with nocturnal hypoxaemia and short burst oxygen for palliation of dyspnoea. This is despite a lack of supporting evidence from randomized clinical trials to support these indications[3-5].

A clinical indication that stimulates long-standing debate is the specific use of supplemental oxygen during exercise training in patients with EID. There is a sound physiological rationale in that supplemental oxygen can reduce respiratory rate and dynamic hyperinflation[6]. It also improves oxygen delivery, leading to a reduction in exercise-associated metabolic acidosis [7]. More than sixty years ago, Cotes and colleagues demonstrated that ambulatory oxygen could support outpatients with chronic respiratory insufficiency during exercise by improving arterial oxygen saturation and increasing walking time[8]. Subsequent, largely laboratory-based, studies have demonstrated a positive, albeit heterogeneous, benefit in exercise time following acute administration of oxygen during an exercise test in patients with COPD or cystic fibrosis[9, 10]. This had led to the logical hypothesis that by improving exercise time acutely, supplemental oxygen allows patients to tolerate training sessions of higher intensity with consequently gain greater benefit from an exercise training programme.

Although there have been attempts to provide evidence to support this hypothesis [11-14], the results from clinical trials have been mixed and inconclusive, largely due to methodological issues, small sample sizes and inclusion of a heterogeneous population, including existing long-term oxygen users [11] or individuals without EID [13, 14].

In this issue of the *European Respiratory Journal*, Alison and colleagues report on a double-blind randomised placebo-controlled trial that aimed to determine whether supplemental oxygen administered during exercise training was more effective than medical air in improving endurance exercise capacity; health-related quality of life; peak walking capacity; dyspnoea and physical activity levels in patients with COPD who were normoxaemic at rest but desaturated during exercise (<90% during the six-minute walk test)[15]. Following random allocation to either oxygen or compressed air, both groups underwent an eight-week training programme with three supervised exercise sessions per week. Training initially consisted of 30 minutes of individualised aerobic exercise (20 minutes of treadmill walking and 10 minutes of stationary cycling). Training intensity was increased according to target dyspnoea scores (3-4 on modified dyspnoea scale [16]) and training duration was increased according to a fixed protocol. The results of this study demonstrated that both groups improved exercise capacity and health-related quality of life following exercise training, but no added benefit was seen with supplemental oxygen in either of the primary outcomes measures of endurance shuttle walk test and chronic respiratory questionnaire, nor in any of the secondary outcome measures. This was a well-designed and conducted study, with clear eligibility criteria, high retention of participants, and successful blinding of participants, trainers and assessors: a feature absent in previous trials[11-14].

A fundamental observation was that the oxygen group were not able to achieve a greater training dose per session than the compressed air group, despite a significantly higher measured oxygen saturation and significantly lower dyspnoea and perceived exertion scores during treadmill training sessions. Without an increased training load, it was not altogether surprising that no between-group differences in response to exercise-training were observed. The authors speculated that the large physiological stimulus applied to both groups overwhelmed the small physiological advantage of acute oxygen administration. Although this is a highly plausible explanation (observed in previous pulmonary rehabilitation adjunct trials[17, 18]), other mechanisms may have limited exercise-training load. The authors did not provide data regarding leg effort or lower limb symptoms, which are reported frequently as a reason for terminating exercise in a proportion of patients with COPD[19]. The exercise-training programme was protocolized such that exercise type and frequency were fixed and exercise duration was progressed up to a maximum of 40 minutes by week three. From this point onwards, staff supervising exercise-training could only increase training intensity (speed of treadmill and cycle workload) but not duration of training – the parameter that is most consistently influenced by acute oxygen response studies. Another speculative explanation is that because the trainers were blinded to treatment allocation and to patients' oxygen saturation during training, they may have inadvertently been more cautious of increasing training intensity in patients known to desaturate during exercise (of whom half were using air). Circumstantial evidence to support this is the observation that the responses to exercise-training in both groups, although statistically significant, were smaller than expected and did not achieve the minimum clinically important difference of the incremental shuttle walk test nor some domains of the Chronic Respiratory Questionnaire following the exercise-training programme.

Another important consideration is that most participants had moderate EID, and the trial was therefore underpowered to detect an effect of supplemental oxygen in patients with more severe EID (nadir below 80%). Furthermore, prior to the start of the exercise programme, the acute physiological response to oxygen was not tested, hence it is not clear as to the proportion of acute oxygen responders and non-responders in the treatment and control groups. Hence the results of the trial cannot be extrapolated to the specific subgroups of patients with COPD and severe EID, those who have shown a beneficial response to acute oxygen administration, nor patients with non-COPD disorders (such as idiopathic pulmonary fibrosis and pulmonary hypertension) and EID. Further research is required to understand the role of supplemental oxygen in the exercise-training of these patient subgroups.


Alison and colleagues should be congratulated in rigorously conducting the largest study to date to examine the role of supplemental oxygen during exercise-training in patients with COPD and EID. They provide convincing data that the majority of patients with COPD and EID safely benefit from exercise-training without the need for supplemental oxygen. This supports the recommendations of international pulmonary rehabilitation guidelines that supplemental oxygen should not be routinely prescribed in patients with COPD who desaturate but do not fulfil indications for long-term oxygen therapy [20-22]. The findings therefore may help to simplify the assessment process of patients with EID for exercise-training interventions such as pulmonary rehabilitation. We present our current assessment algorithm prior to commencement of pulmonary rehabilitation (Figure 1), which acknowledges remaining uncertainty around the role of supplemental oxygen in specific patient subgroups, namely those who show an acute response to supplemental oxygen and those with severe EID.

Although Alison and colleagues intended to explore whether exercise-training or pulmonary rehabilitation in patients with EID could be enhanced with supplemental oxygen, perhaps the most important clinical implication of this study is that for the vast majority of patients, the non-availability of supplemental oxygen need not be a barrier to using community and non-medical settings to deliver pulmonary rehabilitation. This potentially increases the accessibility, utilization and reach of an evidence-based intervention that is considerably more effective in reducing symptom burden in COPD than supplemental oxygen alone.

## Figure Legend

Figure 1: An algorithm to assessing exercise induced oxygen desaturation (EID) (defined as nadir oxygen saturation below 90% during a field walking test performed on room air) in patients with COPD referred for pulmonary rehabilitation.





## References:

1. *Long term domiciliary oxygen therapy in chronic hypoxic cor pulmonale complicating chronic bronchitis and emphysema. Report of the Medical Research Council Working Party.* Lancet 1981; 1(8222): 681-6.
2. Nocturnal Oxygen Therapy Trial G, *Continuous or Nocturnal Oxygen Therapy in Hypoxemic Chronic Obstructive Lung Disease: A Clinical Trial.* Annals of Internal Medicine 1980; 93(3): 391-398.
3. Ameer F, Carson KV, Usmani ZA, Smith BJ, *Ambulatory oxygen for people with chronic obstructive pulmonary disease who are not hypoxaemic at rest.* Cochrane Database of Systematic Reviews 2014(6).
4. Chaouat A, Weitzenblum E, Kessler R, Charpentier C, Enrhart M, Schott R, Levi-Valensi P, Zielinski J, Delaunois L, Cornudella R, Moutinho dos Santos J, *A randomized trial of nocturnal oxygen therapy in chronic obstructive pulmonary disease patients.* Eur Respir J 1999; 14(5): 1002-8.
5. Uronis HE, Ekstrom MP, Currow DC, McCrory DC, Samsa GP, Abernethy AP, *Oxygen for relief of dyspnoea in people with chronic obstructive pulmonary disease who would not qualify for home oxygen: a systematic review and meta-analysis.* Thorax 2015; 70(5): 492-4.
6. Somfay A, Porszasz J, Lee SM, Casaburi R, *Dose-response effect of oxygen on hyperinflation and exercise endurance in nonhypoxaemic COPD patients.* Eur Respir J 2001; 18(1): 77-84.
7. O'Donnell DE, Bain DJ, Webb KA, *Factors contributing to relief of exertional breathlessness during hyperoxia in chronic airflow limitation.* American Journal of Respiratory and Critical Care Medicine 1997; 155(2): 530-535.
8. Cotes JE, Gilson JC, *Effect of oxygen on exercise ability in chronic respiratory insufficiency; use of portable apparatus.* Lancet 1956; 270(6928): 872-6.
9. Bradley JM, O'Neill B, *Short-term ambulatory oxygen for chronic obstructive pulmonary disease.* Cochrane Database Syst Rev 2005(4): Cd004356.
10. Marcus CL, Bader D, Stabile MW, Wang CI, Osher AB, Keens TG, *Supplemental oxygen and exercise performance in patients with cystic fibrosis with severe pulmonary disease.* Chest 1992; 101(1): 52-7.
11. Garrod R, Paul EA, Wedzicha JA, *Supplemental oxygen during pulmonary rehabilitation in patients with COPD with exercise hypoxaemia.* Thorax 2000; 55(7): 539-43.
12. Rooyackers JM, Dekhuijzen PN, Van Herwaarden CL, Folgering HT, *Training with supplemental oxygen in patients with COPD and hypoxaemia at peak exercise.* Eur Respir J 1997; 10(6): 1278-84.
13. Emtner M, Porszasz J, Burns M, Somfay A, Casaburi R, *Benefits of Supplemental Oxygen in Exercise Training in Nonhypoxemic Chronic Obstructive Pulmonary Disease Patients.* American Journal of Respiratory and Critical Care Medicine 2003; 168(9): 1034-1042.
14. Spielmanns M, Fuchs-Bergsma C, Winkler A, Fox G, Kruger S, Baum K, *Effects of Oxygen Supply During Training on Subjects With COPD Who Are Normoxemic at Rest and During Exercise: A Blinded Randomized Controlled Trial.* Respir Care 2015; 60(4): 540-8.
15. Alison JA, McKeough ZJ, Leung RWM, Holland AE, Hill K, Morris NR, Jenkins S, Spencer LM, Hill CJ, Lee AL, Seale H, Cecins N, McDonald CF, *Oxygen compared to air during exercise training in COPD with exercise-induced desaturation.* European Respiratory Journal 2019; 1802429.
16. Borg GA, *Psychophysical bases of perceived exertion.* Med Sci Sports Exerc 1982; 14(5): 377-81.

17. Nolan CM, Maddocks M, Canavan JL, Jones SE, Delogu V, Kaliaraju D, Banya W, Kon SSC, Polkey MI, Man WD, *Pedometer Step Count Targets during Pulmonary Rehabilitation in Chronic Obstructive Pulmonary Disease. A Randomized Controlled Trial*. Am J Respir Crit Care Med 2017; 195(10): 1344-1352.
18. Curtis KJ, Meyrick VM, Mehta B, Haji GS, Li K, Montgomery H, Man WD, Polkey MI, Hopkinson NS, *Angiotensin-Converting Enzyme Inhibition as an Adjunct to Pulmonary Rehabilitation in Chronic Obstructive Pulmonary Disease*. Am J Respir Crit Care Med 2016; 194(11): 1349-1357.
19. Man WD, Soliman MG, Gearing J, Radford SG, Rafferty GF, Gray BJ, Polkey MI, Moxham J, *Symptoms and quadriceps fatigability after walking and cycling in chronic obstructive pulmonary disease*. Am J Respir Crit Care Med 2003; 168(5): 562-7.
20. Spruit MA, Singh SJ, Garvey C, ZuWallack R, Nici L, Rochester C, Hill K, Holland AE, Lareau SC, Man WD-C, Pitta F, Sewell L, Raskin J, Bourbeau J, Crouch R, Franssen FME, Casaburi R, Vercoulen JH, Vogiatzis I, Gosselink R, Clini EM, Effing TW, Maltais F, Palen Jvd, Troosters T, Janssen DJA, Collins E, Garcia-Aymerich J, Brooks D, Fahy BF, Puhan MA, Hoogendoorn M, Garrod R, Schols AMWJ, Carlin B, Benzo R, Meek P, Morgan M, Mølken MPMHR-v, Ries AL, Make B, Goldstein RS, Dowson CA, Brozek JL, Donner CF, Wouters EFM, *An Official American Thoracic Society/European Respiratory Society Statement: Key Concepts and Advances in Pulmonary Rehabilitation*. American Journal of Respiratory and Critical Care Medicine 2013; 188(8): e13-e64.
21. Bolton CE, Bevan-Smith EF, Blakey JD, Crowe P, Elkin SL, Garrod R, Greening NJ, Heslop K, Hull JH, Man WD-C, Morgan MD, Proud D, Roberts CM, Sewell L, Singh SJ, Walker PP, Walmsley S, *British Thoracic Society guideline on pulmonary rehabilitation in adults: accredited by NICE*. Thorax 2013; 68(Suppl 2): ii1-ii30.
22. Alison JA, McKeough ZJ, Johnston K, McNamara RJ, Spencer LM, Jenkins SC, Hill CJ, McDonald VM, Frith P, Cafarella P, Brooke M, Cameron-Tucker HL, Candy S, Cecins N, Chan AS, Dale MT, Dowman LM, Granger C, Halloran S, Jung P, Lee AL, Leung R, Matulick T, Osadnik C, Roberts M, Walsh J, Wootton S, Holland AE, *Australian and New Zealand Pulmonary Rehabilitation Guidelines*. Respiriology 2017; 22(4): 800-819.